

Concise debate on the consequence of the Petroleum Ether mine of the Alovevera Gel in the PCOS mold

Radha Maharjan
Department of Biochemistry,
M.S. University of Baroda, Vadodara, Gujarat, India.

ABSTRACT

This study is related to the consequence of the Petroleum Ether mine of the Alovevera Gel in the PCOS mold. Phytosterols (plant sterols) are secondary plant metabolites which structural and biological counterparts of cholesterol. Plant sterols are responsible for permeability and fluidity of cell membranes. thereby reducing their absorption Phytosterols can be hydrogenated to obtain phyto-stanols. Both esters are chemically stable materials, having comparable chemical and physical properties to edible fats and oils. The substances are insoluble in water, but soluble in non-polar solvents, suchas hexane, iso-octane and 2-propanol.

The research has focused on β sitosterol for the development of steroidal drugs and functional food ingredients among all phytosterols as it exerts hypo-cholesterolemic effects by inhibiting the absorption of cholesterol in the intestines through competition with LDL-cholesterol (Gylling and Miettinen 1996) and also decreases in total, along with LDL-cholesterol concentrations of 5–15% as evident from several studies (Jones et al. 1999; Ostlund Jr 2004). *Aloe vera* gel contains various phytosterols wherein β -sitosterol, stigmasterol, lupeol and campesterol are in abundance (Hamman 2008).

In clinical study, esterified sterol and stanol mixtures consumption resulted in reduction in plasma Triglycerides (TG) concentrations (Jones et al. 2000). This could be correlated with hypoglycaemic effect rendered by AVG. As PCOS a disease condition wherein hyperglycaemia and dyslipidemia are secondary manifestations in addition to ovarian dysfunction. Thereby, these phyto-components independently or synergistically could act at various targets as above mentioned mechanisms. Steroidogenesis is functional attribute of ovary which is under the regulation of hypothalamus-pituitary axis. “*Ex vivo*” study from previous chapter showed wherein non-polar P1 fraction extracted by petroleum ether indicated that it could directly modulate ovarian steroidogenic enzyme activity. Thereby, this research paper attempts to understand “*in vivo*” role of non polar P1 fraction from AVG on steroidogenesis with special reference to ovary.

Key word: petroleum ether, *Aloe vera* gel, Plant sterols, extract, steroidogenesis.

INTRODUCTION

Several reports have suggested various efficacies such as hypoglycaemic, antitumor, anti-inflammatory, antioxidant and laxative effects of *Aloe vera* gel (Rajasekaran et al. 2004; Tanaka et al. 2006). These beneficial properties of *Aloe* leaf have been attributed to the various phyto-compounds found in the inner leaf parenchymatous tissue (Boudreau and Beland 2006; Hamman 2008). The presence of sterols, phenols, alkaloids, polyphenols, flavonoids and lignin have attributed to various efficacies (Madhavi et al. 1995; Veerapur et al. 2009). Recently, several researchers have tried to check in the efficacy of phyto-components from AVG in the content of its hypoglycemic and anti-inflammatory potential (Rajasekaran et al. 2006; Saleem 2009).

One of the well studied phyto-component in *Aloe vera* gel is phytosterol, which has various biological properties (Jones et al. 1999; Ostlund Jr 2004). Phytosterols (plant sterols) are secondary plant metabolites which structural and biological counterparts of cholesterol. Plant sterols are responsible for permeability and fluidity of cell membranes. Further, they act as precursors of brassinosteroids, thus regulating storage and transport processes, transform into numerous other metabolites such as glycoalkaloids and saponins (Piironen et al. 2000). Similarly to cholesterol, most phytosterols have a steroid nucleus and a hydroxyl group in the 3 β -position at carbon atom. Structural differences with cholesterol are mainly found inside chain substitution at C24, an additional double bond between C22 and C23 and/or, in the case of plant stanols, saturation taken place at steroid nucleus. The addition of a methyl/ethyl group at the C-24 position of the side chain sterol (campesterol/ β - sitosterol, respectively) whereas in the case of plant stanols, the saturation of the Δ^5 double bond (campestanol and sitostanol) increases the hydrophobicity of the molecule, thereby reducing their absorption (Heinemann and Ankenbauer 1993). Phytosterols can be hydrogenated to obtain phyto-stanols. Both esters are chemically stable materials, having comparable chemical and physical properties to edible fats and oils. The substances are insoluble in water, but soluble in non-polar solvents, such as hexane, iso-octane and 2-propanol.

An important co-morbidity of hyperglycemia is Dyslipidemia. Thereby, research has focused on β sitosterol for the development of steroidal drugs and functional food ingredients among all phytosterols as it exerts hypo-cholesterolemic effects by inhibiting the absorption of cholesterol in the intestines through competition with LDL-cholesterol (Gylling and Miettinen 1996) and also decreases in total, along with LDL-cholesterol concentrations of 5–15% as evident

from several studies (Jones et al. 1999; Ostlund Jr 2004). *Aloe vera* gel contains various phytosterols wherein β -sitosterol, stigmasterol, lupeol and campesterol are in abundance (Hamman 2008). Also, earlier chapter demonstrated that 70% of P1 fraction (Petroleum ether extracted fraction) is enriched with β sitosterol and other phytosterols. These phytosterols present in *Aloe vera* gel is known to have cholesterol lowering property wherein “*in vitro*” study has demonstrated that phytosterols activates liver X receptor [a nuclear receptor- controls the expression of ATP-binding cassette sub-family G member 5 (ABCG5/8)], thereby increasing the expression of ATP Binding Cassette (ABC) transporters and the transport of cholesterol from enterocytes back into the intestinal lumen, which in turn leads to reduction in cholesterol absorption from intestinal wall (Plat and Mensink 2005).

In clinical study, esterified sterol and stanol mixtures consumption resulted in reduction in plasma Triglycerides (TG) concentrations (Jones et al. 2000). This could be correlated with hypoglycaemic effect rendered by AVG. Also, molecular level studies have shown that decrease in the expression levels of hepatic genes encoding gluconeogenic enzymes [Glucose 6-phosphatase, Phospho-enolpyruvate carboxykinase (PEPCK)], lipogenic enzymes (ACC, FAS), and sterol regulatory element-binding proteins 1 (SREBPs 1) whereas increased the expression of hepatic β -oxidation enzymes and Peroxisome Proliferator Activated Receptor alpha (PPAR α) expressions could alter dyslipidemia.

As PCOS a disease condition wherein hyperglycaemia and dyslipidemia are secondary manifestations in addition to ovarian dysfunction. Also, it is clear that AVG has enriched β sitosterol, stigmasterol, lupeol and fatty acids derivatives as shown in the current study. Thereby, these phyto-components independently or synergistically could act at various targets as above mentioned mechanisms. Steroidogenesis is functional attribute of ovary which is under the regulation of hypothalamus-pituitary axis. “*Ex vivo*” study from previous chapter showed wherein non-polar P1 fraction extracted by petroleum ether indicated that it could directly modulate ovarian steroidogenic enzyme activity. Thereby, current chapter attempts to understand “*in vivo*” role of non polar P1 fraction from AVG on steroidogenesis with special reference to ovary.

RESULTS

The phytochemical analysis of *Aloe vera* gel extracted with petroleum ether exhibited various phytosterols in non-polar fraction has been named as (P1) fraction. Also, “*ex-vivo*” study suggested that P1 fraction has direct effect on ovarian steroidogenic enzyme activity. Hence, in

view of this, aim of present chapter was to evaluate the “*in vivo*” effect of P1 fraction of *Aloe vera* gel in letrozole induced PCOS rat model.

In this context, letrozole induced PCOS rats (P group) were treated with P1 fraction using olive oil as vehicle (~25µg/orally) for 60 days (P1P group). In addition to this, there were PCOS animals which received olive oil and served as vehicle group (VP) while control group (C group) received P1 fraction was termed as herbal control (P1C group).

Data from previous chapters have demonstrated that 10 mg dry weight of *Aloe vera* gel for 60 days was the minimum effective dose required for the management of clinical pathophysiological symptoms of PCOS. In addition to this, a time-dependent profiling of phytocomponents present in the P1 fraction was performed by HPTLC to check their stability upon storage. Results demonstrate that the P1 extract was stable upto 60 days (Figure). Hence, the P1 treatment was continued for 60 days.




Figure: Effect of P1 fraction of AVG on Body weight in letrozole induced PCOS rats

There was no significant alteration in the body weight of animals after treatment with P1 fraction of *Aloe vera* gel (Figure).

** $p < 0.01$, *** $p < 0.001$) at 60, 90 and 120 minutes of the test respectively as compared to control group. At 30 days of time interval, P1 fraction treated PCOS rats (P1P) demonstrated a

significant change at 60, 90 and 120 minutes point of OGTT profile (@ $p < 0.05$, @@ $p < 0.01$, @@@ $p < 0.001$) whereas PCOS rats treated with olive oil (VP) showed significant change at 90 and 120 minute (@ $p < 0.05$) in OGTT profile as compared to PCOS group. Herbal control group (P1C) depicted similar results to control rats (Figure 7.3.3 (A)). At end of experiment regime (60 days), OGTT profile was evaluated again in all groups of animals to check glucose tolerance. Letrozole induced PCOS rats (P).

Table : Effect of P1 fraction of AVG on insulin status in letrozole induced PCOS rats

| GROUPS | Insulin (IU/ml) | HOMA-IR |
|--------|-----------------|----------------|
| C | 12.3±0.6 | 1.8±0.1 |
| P1C | 13.2±1.2 | 2.5±0.1 |
| P | 21±1.0** | 4.5±0.3*** |
| VP | @@@ 9.1±1.2 | @@@ 1.4±0.1 |
| P1P | @ 13.5±0.76 | @@ 2.5±0.2 |

N=4, The values represented as Mean±SEMC=Control; P C= P treated control; P=PCOS; VP=Vehicle (Olive oil) treated PCOS

During the course of experiment, liver toxicity markers like Serum Glutamate Pyruvate Transaminase (SGPT) and creatinine was evaluated. P1 fraction treated groups (P1P) exhibited non-significant change in levels of the toxicity markers. In addition to this, letrozole treated PCOS group (P) as well as olive oil-vehicle treated animals (VP) demonstrated no change in the above parameters upon treatment indicating no toxic effects of phytosterols fraction on PCOS rats during study (Figure 7.3.4).

This hypertriglyceridemia has been normalized with P1 fraction (P1P) treatment (@@@ $p < 0.001$). P1 fraction demonstrated more significant change as compared to Vehicle group (VP) group (# $p < 0.05$) suggested that P1 fraction rich in various phytosterols has more lipid lowering efficacy. In PCOS rats, LDL levels were also reduced after P1 fraction treatment (@@@ $p < 0.001$) whereas no significant changes were observed in HDL and VLDL levels. Vehicle group (VP) also showed significant change in LDL level (@@@ $p < 0.01$) but P1 fraction again exerted more significant change as compared to vehicle (VP) group (### $p < 0.001$) (Table 7.3.2).


Table: Effect of P1 fraction on Lipid profile in letrozole induced PCOS rats

| | Triglycerides | Total Cholesterol | LDL | HDL | VLDL |
|------------|--------------------------|-------------------|----------------------------|----------|-----------|
| C | 70.8±2.9 | 48.3±1.7 | 51.9±3.6 | 31.1±2.0 | 13.8±0.69 |
| P1C | 71.6±4.9 | 52.6±5.7 | 49.6±3.1 | 27.6±2.8 | 16.0±1.5 |
| P | 96.0±6.3*** | 55.8±3.4 | 108.0±10.2*** | 24.5±2.3 | 22.6±2.0 |
| VP | 78.7±2.3 [®] | 43.6±3.7 | 79.11±5.5 ^{®®®} | 34.1±1.4 | 13.7±1.3 |
| P1P | 64.2±2.4 ^{®®®#} | 45.3±4.3 | 45.8±2.2 ^{®®®###} | 28.7±1.4 | 13.2±0.4 |

N=4, The values represented as Mean±SEMC=Control; P C= P treated control; P=PCOS; VP=Vehicle (Olive oil) treated PCOS;

Steroidogenesis plays a major role in regulation of the ovarian functioning and PCO condition is characterized by altered steroidogenesis leading to impairment in folliculogenesis (Diamanti-Kandarakis 2008). Thereby, key steroidogenic enzymes- 3β hydroxysteroid dehydrogenase (3β HSD) and 17β hydroxysteroid dehydrogenase (17β HSD) were studied in reproductive tissues: ovary, uterus and brain tissues: hypothalamus, pituitary in all groups of animals.

Figure: Effect of P1 fraction of AVG on Ovarian histological changes in letrozole



C=Control; P₁C= P₁ treated control; P=PCOS; VP=Vehicle (Olive oil) treated PCOS; P₁P=P₁ treated PCOS
N=3, All sections taken in diestrus stage. Magnification- 4X

▲ : Growing follicles ▲ : Cyst

Ovarian steroid hormones perform several important actions related to reproductive organ function and their effects are mediated through interaction with specific proteins (Drummond and Findlay 1999; Salvetti et al. 2010). Several studies have shown that PCO affects the expression of steroidogenic proteins such as aromatase, 3 β -HSD, LHR and StAR in the ovaries of letrozole treated rats (Zurvarra et al. 2009). Thereby, transcripts levels were evaluated in all groups of animals. Figure elucidated the effect on P1 fraction on transcript level of key protein in steroidogenesis wherein PCOS rat demonstrated a elevated gene expression of StAR (**p<0.001), insulin receptor (IR) (**p<0.01), Luteinizing hormone receptor (**p<0.01) and androgen receptor (AR) (**p<0.001) whereas decrease was observed in gene expression of aromatase (*p<0.05) as compared to control group. However, there was no significant change was observed in gene expression of follicle stimulating hormone receptor (FSHR) in PCOS (P) group. After P1 treatment, there were significant changes observed in these key steroidogenic protein wherein substantial decrease in StAR (**p<0.001), insulin receptor (IR) (*p<0.05), LHR (**p<0.01) and AR (**p<0.01) transcripts level as compared to PCOS rats.5) whereas significant increase was observed in transcript level of aromatase (**p<0.001) in P1 fraction treatment (P1P). However, no significant changes were observed in all transcript level in vehicle (VP) group. Also, P1 fraction treated PCOS rats (P1P) demonstrated more significant change in gene expression of above key protein – androgen receptor (**p<0.01), StAR (**p<0.001), (**p<0.001) as compared to vehicle (VP) group suggested that P1 fraction has more efficacy in comparative study (Figure 7.3.7)

DISCUSSION

Aloe species are now considered a very interesting source of bioactive compounds amongst which phytosterols play a major role (Tanaka et al. 2006; Misawa et al. 2012). Present study was an attempt to investigate the efficacy of partially purified P1 fraction (NPP) of *Aloe vera* gel with impact on modulation of ovarian structure-function in letrozole induced PCOS rat model. Our earlier published data suggested that treatment of *Aloe vera* gel restored glucose intolerance and dyslipidemia along with cyclicity. It has also modulatory effect on ovarian steroidogenesis (Maharjan et al. 2010; Desai et al. 2012).

In view of this result, current chapter investigates “*in vivo*” effect of partially purified non polar fraction of *Aloe vera* gel on letrozole induced PCOS rats. During experimental regime, stability of P1 fraction of *Aloe vera* gel as various phyto- components (volatile in nature) present

in gel was assayed. HPTLC data suggested phytosterols are stable up to 60 days (Time period of *in vivo* treatment being 60 days). During course of experiments, levels of toxicity markers were estimated wherein no significant changes were observed in Serum Glutamate Pyruvate Transaminase (SGPT) and creatinine level. This may be because of low dose of phytosterols (~25 µg) that did not exert any toxic effects in PCOS rats.

Letrozole induced PCOS positive rats' demonstrated PCO phenotype including irregular cyclicality, increased body weight, glucose intolerance and altered steroid hormones profile. After validation of PCOS rat model, these PCOS positive rats were treated with partially purified non polar P1 fraction of *Aloe vera* gel (~25µg/orally) for 60 days. P1 fraction treated PCOS rats exhibited restored the glucose sensitivity as mediated by improved HOMA-IR implicating that phytosterols of P1 has good glucose lowering property (Tanaka et al. 2006). There are various reports that suggesting anti-hyperglycaemic property of phytosterols that is identified from *Aloe vera* gel which improved glucose intolerance (Tanaka et al. 2006; Misawa et al. 2008)) has also demonstrated that sterols isolated from *Aloe vera* reduced the expression of gluconeogenic enzymes (G6 Pase and PEPCK) and also stimulated glucose catabolism along with suppressed *de novo* glucose production. Also, phytosterols have direct effect on glucose metabolism wherein synergism of β- sitosterol and stigmasterol produced hypoglycaemic activity (Jamaluddin et al. 1994).

It is known hyperglycemia leads to hyperinsulinemia. Also, it considered as one of the main complication that add to metabolic complexity in PCO phenotype. Present study showed that PCOS rats with elevated insulin level exhibited high HOMA-IR index suggesting hyperinsulinemia with insulin resistance. Also, insulin receptor up-regulation is noted in PCOS phenotype. This further strengthens that letrozole induced PCOS rats are insulin resistant. The values returned to normal level when treated with P1 fraction (phytosterols rich) suggesting that it has potential to sensitize the insulin receptor and reduce insulin level in PCO condition; thereby reverting insulin resistant state to sensitive status as indicated by improved HOMA-IR change. This could be evident from that reports which indicate the potentiality of phytosterols of AVG as insulin sensitizer (Misawa et al. 2012). Phytosterols isolated from different plants also reported to have anti hyperglycaemic effect and improved insulin sensitivity (Russell et al. 2002; Tanaka et al. 2006).

Apart from glucose metabolism, PCOS rats also exhibited elevated triglycerides and LDL

level which are important contributors for lipid metabolism (Hussain and Alam 2014). In present study, dyslipidemia was again restored after treatment of P1 fraction indicating lipid lowering property of P1 fraction. Clinical study also suggested that consumption of phytosterol- β -sitosterol helps to decrease plasma low density lipoprotein (LDL) and triglyceride (TG) levels; although there was usually no change was observed in high-density lipoprotein (HDL) (Jones et al. 2000). In this context, phytosterols isolated from AVG is known to have direct effect on lipogenic enzymes that involved in lipid biosynthesis wherein these isolated phytosterols administration induces down regulation of fatty acid synthesis and a tendency for up-regulation of fatty acid oxidation in the liver that lead to reduction in intra-abdominal fat and improvement of hyperlipidemia (Misawa et al. 2012). Lupeol, a antioxidant rich compound present in P1 fraction also has properties to normalize the lipid profile by decreasing LDL and total cholesterol level along with improved antioxidant status (Sudhahar et al. 2006). P1 fraction containing β - sitosterol, has a beneficial role in the reduction of the intestinal uptake of cholesterol and lowering the concentration of cholesterol in dietary mixed micelles via a dynamic competition mechanism (Mel'nikov et al. 2004). Our study has also similar kind of alteration which may due to synergistic action of phytosterols present in AVG. However, role of each phyto-components present in P1 need to be studied further.

In this context, regulation of steroidogenesis governed by level of gonadotropin receptors and steroid receptors expressed on ovarian cells wherein expression of LHR and Androgen receptor is shown to be elevated on the stimulation by high insulin level in PCOS rats which further lead to high theca androgen modulation- Testosterone, DHEA, androstenedione. This may be due to high 3β -HSD enzyme activity, which is attributed by increased expression of transcript and protein. Also, heightened androgen receptor (AR) expression further increases the androgen production through 3β -HSD activity. Also, it has been demonstrated that insulin acts directly via its own insulin receptor at physiological concentrations in cultured polycystic ovary theca cells (Nestler 1997; Nelson-Degrave et al. 2005) and increases androgen production (Nelson et al. 1999), which is significantly greater in ovarian theca cells than from normal women. Thereby, increased expression level of insulin receptor (IR) in PCO rats could also contribute to rise in androgen content as seen in present study. Insulin alone can also increase androstenedione production and can synergize with LH to increase androgen biosynthesis in PCO condition (Nelson et al. 1999). This elevated IR expression was reduced when treated with P1 fraction of AVG. This may be due to phytosterols present in P1 fraction that can act as insulin receptor

sensitizer in diabetic rat (Tanaka et al. 2006).

In addition to this, LH also can upregulate StAR expression via cAMP signalling in ovary by binding with its receptor LHR present on the ovaries (Bauer et al. 2000). Elevated expression of LHR was noted in the current study. This could be due to increase serum LH concentrations, as reported by many scientists (Maliqueo et al. 2012; Kauffman et al. 2015). Transcript level of LHR was decreased while treated with P1 fraction of AVG. This may be because of the estrogenic effects of β -sitosterol on the pituitary wherein studies of phytoestrogens in mammals have demonstrated that genistein decreases GnRH-induced luteinizing hormone (LH) in rats (Hughes et al. 2004) and that coumesterol (phytosterol) decreases the amplitude of LH pulses in ewes (MacLatchy and Vanderkraak 1995). Also, β -Sitosterol reduces the rate of the side chain cleavage reaction, as compared to cholesterol activity (Masuo et al. 1980). Decreased pregnenolone production was concomitant with decrease in T production (MacLatchy and Vanderkraak 1995). This suggests that β -sitosterol could be directly affecting steroidogenic enzymes within the biosynthetic pathway.

With reference to above context, aromatase also plays a crucial role in steroid status wherein it converts theca androgens into estrogens, which have role in follicular development and regulation of feedback mechanism of reproductive cycle. In present study, transcript level as well as relative protein expression level was decreased in PCOS rats suggesting disturbed testosterone: estrogen flux due to lower conversion and thus leading to excess androgen production though ovarian steroidogenesis. Also, decreased aromatase expression is well correlated with low level of estradiol as seen in PCOS rats. However, altered expression level observed in PCO phenotype was reverted back to normal when treated with P1 fraction of AVG. This may be attributed to the fact that StAR protein has been normalized by phytosterols, which reorients its flux where androgen is converted into estrogens efficiently by aromatase. Also, it is to be noted P1 components independently have potential to up-regulate the steroidogenic proteins as evident from P1 group.

Any functional modulation is mainly reflection of structural aberrations. Structurally, PCO rats demonstrated the presence of multiple fluid filled peripheral cysts in the ovary. Several studies have shown similar results (Kafali et al. 2004). P1 fraction treated PCOS rats demonstrated a reduction in peripheral cysts in ovary and increased growing follicles with presence of corpus luetum that mainly indicates successful ovulation and release of dominant mature graffian follicle.

The restoration in the ovarian structure and function can be attributed to phytosterols that lead to modulation in the HPO axis. This modulation helped in maturation of follicles and release matured ova during ovulation. Reports have suggested that various medicinal ayurvedic herbs that helped to restore the ovulation and fertility (Adimoelja 2000; Sasikala et al. 2010). However, role of individual phytosterol at molecular level needsto be studied.

CONCLUSION

Current chapter focused on evaluation of “*in vivo*” efficacy of non polar P1 fraction of *Aloe vera* gel in letrozole induced PCOS rat model wherein P1 fraction (enriched with β sitosterol, stigmasterol, lupeol as major phytosterols) treated PCOS rats demonstrated glucose sensitivity and regular cyclicity. It also restored lipid profile and disturbed hormonal profile that again helped to repair ovarian dysfunction in PCO phenotype. Apart from organ level, P1 fraction of *Aloe vera* gel also significantly affects transcripts level of StAR, LHR, androgen receptor (AR), aromatase and Insulin receptor (IR) as well as relative protein expression of StAR, 3β HSD and aromatase in PCOS rats, which may be one of possible reason for modulated steroidogenesis. This implies that P1 fraction rich in β sitosterol, stigmasterol, lupeol and other sterol derivatives that has potential to act at various targets for restoration of ovarian function in PCO phenotype. Overall, study implies that non polar P1 fraction containing phytosterols that could be active phyto- component involved in modulation of hyperglycaemic, hyperinsulinemic condition leading to functional ovulation. These phytocomponents might help us to design future drug that can improve ovarian structure-function. **Overall, studies prove that phytosterols containing P1 fraction of *Aloe vera* gel could be possible novel drug in management of PCOS.**

REFERENCES

1. Awad AB, Fink CS. 2000. Phytosterols as anticancer dietary components: evidence and mechanism of action. *The Journal of nutrition* **130**: 2127-2130.
2. Batta AK, Xu G, Honda A, Miyazaki T, Salen G. 2006. Stigmasterol reduces plasma cholesterol levels and inhibits hepatic synthesis and intestinal absorption in the rat. *Metabolism* **55**: 292-299.
3. Carter BA, Taylor OA, Prendergast DR, Zimmerman TL, Von Furstenberg R, Moore DD, Karpen SJ. 2007. Stigmasterol, a soy lipid-derived phytosterol, is an antagonist of the bile acid nuclear receptor FXR. *Pediatric research* **62**: 301-306.

4. Desai BN, Maharjan RH, Nampoothiri LP. 2012. Aloe barbadensis Mill. formulation restores lipid profile to normal in a letrozole-induced polycystic ovarian syndrome rat model. *Pharmacognosy research* **4**: 109.
5. Gupta M, Kiran S, Gulati A, Singh B, Tewari R. 2012. Isolation and identification of phosphate solubilizing bacteria able to enhance the growth and aloin-A biosynthesis of Aloe barbadensis Miller. *Microbiological research* **167**: 358-363.
6. Gylling H, Miettinen TA. 1996. Effects of inhibiting cholesterol absorption and synthesis on cholesterol and lipoprotein metabolism in hypercholesterolemic non-insulin-dependent diabetic men. *Journal of lipid research* **37**: 1776-1785.
7. Hughes E, Collins J, Vandekerckhove P. 2004. Gonadotrophin-releasing hormone analogue as an adjunct to gonadotropin therapy for clomiphene-resistant polycystic ovarian syndrome. *The Cochrane Library*.
8. Huseini HF, Kianbakht S, Hajiaghaee R, Dabaghian FH. 2012. Anti-hyperglycemic and anti- hypercholesterolemic effects of Aloe vera leaf gel in hyperlipidemic type 2 diabetic patients: a randomized double-blind placebo-controlled clinical trial. *Planta Medica-Natural Products and Medicinal Plant Research* **78**: 311.
9. Kafali H, Iriadam M, Ozardalı I, Demir N. 2004. Letrozole-induced polycystic ovaries in the rat: a new model for cystic ovarian disease. *Archives of medical research* **35**: 103-108.
10. Maharjan R, Nagar PS, Nampoothiri L. 2010. Effect of Aloe barbadensis Mill. formulation on Letrozole induced polycystic ovarian syndrome rat model. *Journal of Ayurveda and integrative medicine* **1**: 273.
11. Nelson-Degrave VL, Wickenheisser JK, Hendricks KL, Asano T, Fujishiro M, Legro RS, Kimball SR,
12. Rajasekaran S, Ravi K, Sivagnanam K, Subramanian S. 2006. Beneficial effects of Aloe vera leaf gel extract on lipid profile status in rats with streptozotocin diabetes. *Clinical and Experimental Pharmacology and Physiology* **33**: 232-237.
13. Russell JC, Ewart HS, Kelly SE, Kralovec J, Wright JL, Dolphin PJ. 2002. Improvement of vascular dysfunction and blood lipids of insulin-resistant rats by a marine oil-based phytosterol compound. *Lipids* **37**: 147-152.
14. Shelton RM. 1991. Aloe vera. *International journal of dermatology* **30**: 679-683.
15. Veerapur V, Prabhakar K, Parihar VK, Kandadi MR, Ramakrishana S, Mishra B, Satish Rao B, Srinivasan K, Priyadarsini K, Unnikrishnan M. 2009. Ficus racemosa stem bark extract: a potent antioxidant and a probable natural radioprotector. *Evidence-Based Complementary and Alternative Medicine* **6**: 317-324.